



ELSEVIER



ScienceDirect

Precision measurement of the $3d \rightarrow 2p$ x-ray energy in kaonic ^4He

S. Okada^{a,*}, G. Beer^b, H. Bhang^c, M. Cargnelli^d, J. Chiba^e, Seonho Choi^c, C. Curceanu^f,
Y. Fukuda^g, T. Hanaki^e, R.S. Hayano^h, M. Iio^a, T. Ishikawa^h, S. Ishimotoⁱ, T. Ishiwatari^d,
K. Itahashi^a, M. Iwaiⁱ, M. Iwasaki^{a,g}, B. Juhász^d, P. Kienle^{d,j}, J. Marton^d, Y. Matsuda^a,
H. Ohnishi^a, H. Oota^a, M. Sato^{g,l}, P. Schmid^d, S. Suzukiⁱ, T. Suzuki^a, H. Tatsuno^h, D. Tomono^a,
E. Widmann^d, T. Yamazaki^{a,h}, H. Yim^c, J. Zmeskal^d

^a *RIKEN Nishina Center, RIKEN, Saitama 351-0198, Japan*

^b *Department of Physics and Astronomy, University of Victoria, British Columbia V8W 3P6, Canada*

^c *Department of Physics, Seoul National University, Seoul 151-742, South Korea*

^d *Stefan Meyer Institut für subatomare Physik, Austrian Academy of Sciences, A-1090 Vienna, Austria*

^e *Department of Physics, Tokyo University of Science, Chiba 278-8510, Japan*

^f *Laboratori Nazionali di Frascati, INFN, I-00044 Frascati, Italy*

^g *Department of Physics, Tokyo Institute of Technology, Tokyo 152-8551, Japan*

^h *Department of Physics, The University of Tokyo, Tokyo 113-0033, Japan*

ⁱ *High Energy Accelerator Research Organization (KEK), Ibaraki 305-0801, Japan*

^j *Physik Department, Technische Universität München, D-85748 Garching, Germany*

Received 30 June 2007; accepted 13 August 2007

Available online 17 August 2007

Editor: V. Metag